

Fermion propagators in QED₃ with velocity anisotropies

Iorwerth Owain Thomas*

University of Wales Swansea

E-mail: pyiori@swansea.ac.uk

Simon Hands

University of Wales Swansea

E-mail: s.hands@swansea.ac.uk

QED₃ with fermi and gap anisotropies is considered to be a candidate effective field theory of high temperature superconductors. Simulations of a variant of the theory have demonstrated that there is evidence consistent with a phase transition to a chirally restored phase as the velocity anisotropy increases, and that the correlation lengths of pions in spatial directions behave in a manner consistent with the anisotropy being a relevant parameter of the theory. We present the first measurements of the fermion propagator in Landau gauge for this theory and suggest that the structure of this theory is more complex than previously thought.

XXIIIrd International Symposium on Lattice Field Theory

25-30 July 2005

Trinity College, Dublin, Ireland

*Speaker.

1. Introduction

QED₃ has been the subject of recent interest in condensed matter theory as a candidate effective theory of the underdoped phases of cuprate high-temperature superconductors [1, 2].

These effective theories have an interesting peculiarity: since we have no reason to suspect that Lorentz invariance is respected in a condensed matter system, we can introduce ‘velocities’ v_F and v_Δ that parametrise the deformation of the Fermi surface of the underlying condensed matter theory. At low energies, our action is [4]:

$$S = \int d^2\mathbf{r} d\tau \{ \bar{\Psi}_1 [\gamma_0(\partial_\tau + ia_\tau) + \frac{\delta}{\sqrt{\kappa}} \gamma_1(\partial_x + ia_x) + \delta \sqrt{\kappa} \gamma_2(\partial_y + ia_y)] \Psi_1 + (1 \rightarrow 2, x \leftrightarrow y) + \frac{1}{4e^2} F_{\mu\nu} F_{\mu\nu} \} \quad (1.1)$$

where Ψ_i is an excitation around one of the two pairs of nodes possessed by the d-wave Fermi surface, $\kappa = \frac{v_F}{v_\Delta}$ (this quantity is proportional to the doping of a cuprate sample), $\delta = \sqrt{v_\Delta v_F}$ and the gauge fields model statistical interactions between excitations at the nodes.

In this approach, it has been suggested that the chirally-broken phase of the model corresponds to the onset of a ‘spin-density wave’ phase [2, 3], and that the chirally symmetric phase to the so-called ‘pseudogap’ phase, some of whose peculiar properties, it is hoped, may be explained by chirally-symmetric QED₃ [1].

In order that we might make use of an exact HMC algorithm, we use the following simplified version of (1.1), which we expect to capture most of the physics of the system:

$$S = \sum_{i=1}^2 \sum_{x,x'} a^3 \bar{\chi}_i(x) M_{x,x'} \chi_i(x') + \frac{\beta}{2} \sum_{x,\mu < \nu} a^3 \Theta_{\mu\nu}^2(x) \quad (1.2)$$

where $M_{x,x'}$ is the staggered fermion matrix, modified as follows,

$$M_{x,x'} = \frac{1}{2a} \sum_{\mu=1}^3 \xi_\mu(x) [\delta_{x',x+\hat{\mu}} U_{x\mu} - \delta_{x',x-\hat{\mu}} U_{x'\mu}^\dagger] + m \delta_{\mu\nu}, \quad (1.3)$$

where $\xi_\mu(x) = \lambda_\mu \eta_\mu(x)$: $\eta_\mu(x) = (-1)^{x_1 + \dots + x_{\mu-1}}$, which is the Kawamoto-Smit phase factor (where $x_1 = x$, $x_2 = y$ and $x_3 = t$), and $\lambda_x = \kappa^{-\frac{1}{2}}$, $\lambda_y = \kappa^{\frac{1}{2}}$, $\lambda_t = 1$.

$U_{x\mu} \equiv \exp(ia\theta_{x\mu})$ is the gauge parallel transporter, and the gauge action has the *non-compact* form

$$\Theta_{\mu\nu}(x) = \frac{1}{a} [\Delta_\mu^+ \theta_\nu(x) - \Delta_\nu^+ \theta_\mu(x)]. \quad (1.4)$$

In [6], we presented evidence from simulations on a 16^3 lattice of a phase transition to what appears to be a chirally symmetric state as κ increases above a critical value, $\kappa_c \approx 4.35$, and that, contrary to previous analytical studies, the renormalised value of κ (when defined as the ratio between the pion screening lengths in the x - and y -directions) is a relevant parameter (i.e. $(\kappa_{\pi ren} - 1) > (\kappa_{bare} - 1)$). In what follows, we present measurements of the fermion propagator in Landau gauge, whose behaviour is thought (not uncontroversially) [1, 5], to provide some explanation for the behaviour of the pseudogap phase.

2. Gauge Fixing

The fermion propagator is not a gauge invariant quantity; prior to measuring it we must numerically impose a gauge fixing condition. We do so as follows [7, 8]:

- We impose ‘minimal’ Landau gauge, corresponding to the continuum condition $\partial_\mu \theta_\mu = 0$:

$$\sum_\mu \theta_{\mu x} - \theta_{\mu x - \hat{\mu}} = 0. \quad (2.1)$$

This corresponds to the minimum of a functional $F[\theta]$. For reasons of practicality, in numerical simulations one imposes the condition up to a residual value, R , which we choose to be the floating point value 10^{-6} .

- To remove all remaining gauge degrees of freedom (largely due to zero modes that are an artefact of the finite volume of the lattice), we rotate $\theta_{\mu x} \mapsto \theta_{\mu x} \pm n \frac{2\pi}{L_\mu}$, $\forall \theta_{\mu x}, \mu$, where n is an integer and L_μ is the size of the lattice in that direction, such that:

$$-\frac{2\pi}{L_\mu} < \bar{\theta}_\mu \leq \frac{2\pi}{L_\mu}, \quad (2.2)$$

where $\bar{\theta}_\mu = \frac{1}{V} \sum_x \theta_{\mu x}$.

We call this the ‘modified non-compact iterative Landau gauge’ or ‘miLandau gauge’ (by analogy with the similarly named condition proposed in [9] for compact QED).

It is, of course, necessary to check that there are no false minima of $F[\theta]$ (Gribov copies) that might contaminate our measurements; while this gauge condition has been used before, this check has yet to be carried out for noncompact QED₃, simply as it seems unlikely that there could be any such problem. It would be good to confirm such an intuition (particularly as compact QED₃ has a non-trivial Gribov problem [10]).

In order to do this, we followed the method of [11]: From a set of mother configurations, we produced 3 sets of 500 daughter configurations via the following gauge transformations:

A: $\theta_{\mu x} \mapsto \theta_{\mu x} - \alpha_{x+\hat{\mu}} + \alpha_x$, where α_x is a random number.

B: $\theta_{\mu x} \mapsto \theta_{\mu x} \pm n \frac{2\pi}{L_\mu}$, $\forall \theta_{\mu x}$ in the direction μ , where n is randomly chosen to be 0, 1, or -1.

C: We perform transformation **A** and then **B**.

We then define a ‘variance’ $dF = \max_{ij} [F[\theta]_i - F[\theta]_j]$, where i, j run over a mother configuration and all its daughters. If, for a given mother and its offspring, $dF < R$, then for all practical purposes, there are no Gribov copies. Our results are presented in Figure 1 for a 16^3 lattice with $m = 0.03$, $\beta = 0.2$ and $\kappa = 1.00$ (we obtained similar results for a value of $\kappa = 10.00$). We see that for set **A**, there are no gauge copies for either gauge condition. However, for **B** and **C**, we see that there are Gribov copies present for mLandau gauge, but not for miLandau gauge. Furthermore, the variance is more-or-less identical for both of these groups, which implies that the source of Gribov copies are rotations of kind found in group **B**. We conclude that, in line with our expectations, the miLandau conditions fix the gauge completely.




Figure 1: Variances for the sets of gauge transformations **A**, **B** and **C**.

3. Fermion Propagator

Fermion propagators for $\beta = 0.2$, $m = 0.03$ on a 16^3 lattice were measured over 30,000 trajectories for $1 \leq \kappa \leq 10$, the extremely large number of trajectories being necessitated by the noisiness of the observable. A very slight tendency towards sawtooth behaviour was noted in the time-direction propagators above κ_c as the value of the propagator on even-numbered slices began to drop; this was more pronounced in the y -direction and absent in the x -direction. It is not clear why this should be; it may have something to do with the restoration of chiral symmetry, though it does not quite replicate the expected behaviour of a fermion propagator under such circumstances ([12] gives an example of what we expect). Ideally, one would like to fit data with this behaviour to a four parameter fit; however, these fits proved to be unstable on this lattice size, and so we fitted a two-parameter curve to the odd timeslices only.

Our results are presented in Figures 2 and 3, where κ_{fr} , the renormalised anisotropy with respect to fermions, is the ratio of the fermion screening mass in the x -direction with that in the y -direction. We may draw two conclusions here: firstly, that fermions retain a dynamically generated mass above κ_c , and secondly that $\kappa_{fren} - 1 < \kappa_{bare} - 1$ – that is, κ is an *irrelevant* parameter with respect to fermions. This last is quite puzzling, given that $\kappa_{\pi ren}$ is a relevant parameter of the theory, and may take some thought to resolve.

This retention of the dynamically generated mass into an apparently chirally-symmetric phase is more readily explicable, however, and has also been observed in the $2 + 1D$ Gross-Neveu model at finite temperature [12]. Witten [13] suggests that such behaviour may be observed if:

- The physical fermion corresponds to a branch-cut, and not a pole, and is constantly interact-




Figure 2: Fermion masses for various values of κ .




Figure 3: The renormalised anisotropy, κ_{fren} , for various κ .

ing with massless, non-Goldstone bosons.

- $\langle \bar{\psi}\psi \rangle$, the chiral condensate, is *close* to having long-range order – for example, it has a phase parameter that slowly varies with spatial/temporal position. This would entail an overall non-zero magnitude of the order parameter, but fermions would still experience a dynamically generated mass.

If correct, this behaviour is unanticipated by any analytic prediction that we are aware of. Because of this, it renders the status of the condensed matter model a little unclear.

Nevertheless, more research is needed before this can be clarified – the thermodynamic and continuum limits must be taken, for example, and it would be good to have some measurements of the fermion mass closer to the chiral limit. Even if this theory proves not to be viable as a model of a condensed matter theory, we hope that its behaviour is interesting enough in its own right to be deserving of study.

4. Acknowledgements

SJH is supported by a PPARC Senior Research Fellowship, IOT by a University of Wales Postgraduate Research Scholarship.

References

- [1] M. Franz, Z. Tešanović and O. Vafek, *QED_3 theory of pairing pseudogap in cuprates I: From d-wave superconductor to antiferromagnet via algebraic Fermi liquid*, *Phys. Rev. B* **66** (2002) 054535 [arXiv:cond-mat/0203333].
- [2] I. Herbut, *QED_3 theory of underdoped high temperature superconductors*, *Phys. Rev. B* **66** (2002) 094504 [arXiv:cond-mat/0202491].
- [3] Z. Tešanović, O Vafek and M Franz, *Chiral Symmetry Breaking and Phase Fluctuations in Cuprate Superconductors: A QED_3 Unified Theory of the Pseudogap State*, *Phys. Rev. B* **65** 180511 [arXiv:cond-mat/0110253].
- [4] D. Lee and I. Herbut, *The effect of velocity anisotropy on the antiferromagnetic instability of a d-wave superconductor*, *Phys. Rev. B* **66** (2002) 094512 [arXiv:cond-mat/0201088].
- [5] D. V. Khveshchenko, *Gauge (non-)invariant Green functions of Dirac fermions coupled to gauge fields*, *Phys. Rev. B* **65** (2002) 235111 [arXiv:cond-mat/0112202];
D. V. Khveshchenko, *Elusive physical electron propagator in QED-like effective theories*, *Nucl. Phys. B* **642** (2002) 515;
V. P. Gusynin, D. V. Khveshchenko and M. Reenders, *Anomalous dimensions of gauge-invariant amplitudes in massless gauge theories*, *Phys. Rev. B* **67** (2003) 115201 [arXiv:cond-mat/0207372].
- [6] S. Hands and I. O. Thomas, *Lattice study of anisotropic QED_3* , *Phys. Rev. B* **72**, 054526 (2005) [arXiv:hep-lat/0412009].
- [7] A. Nakamura and R. Sinclair, *Fermion Propagators In $U(1)$ Lattice Gauge Theory*, *Phys. Lett. B* **243** (1990) 396.
- [8] M. Gockeler, R. Horsley, E. Laermann, U. J. Wiese, P. Rakow, G. Schierholz and R. Sommer, *The Continuum Limit Of QED: Renormalization Group Analysis And The Question Of Triviality*, *Phys. Lett. B* **251** (1990) 567 [Erratum-ibid. *B* **256** (1991) 562].
- [9] S. Durr and P. de Forcrand, *Modified iterative versus Laplacian Landau gauge in compact $U(1)$ theory*, *Phys. Rev. D* **66** (2002) 094504 [arXiv:hep-lat/0206022].
- [10] V. K. Mitrjushkin, *Classical solutions in lattice gauge theories*, *Phys. Lett. B* **389** (1996) 713 [arXiv:hep-lat/9607069];
V. K. Mitrjushkin, *Gauge fixing, zero-momentum modes and the calculation of masses on a lattice*, *Phys. Lett. B* **390** (1997) 293 [arXiv:hep-lat/9606014].
- [11] I. L. Bogolubsky, V. K. Mitrjushkin, M. Muller-Preussker and P. Peter, *Lorentz gauge and Gribov ambiguity in the compact lattice $U(1)$ theory*, *Phys. Lett. B* **458** (1999) 102 [arXiv:hep-lat/9904001].
- [12] S. J. Hands, J. B. Kogut and C. G. Strouthos, *The $(2+1)$ -dimensional Gross-Neveu model with a $U(1)$ chiral symmetry at non-zero temperature*, *Phys. Lett. B* **515** (2001) 407 [arXiv:hep-lat/0107004].
- [13] E. Witten, *Chiral Symmetry, The $1/N$ Expansion, And The $SU(N)$ Thirring Model*, *Nucl. Phys. B* **145** (1978) 110.